4200 Rocklin Road, Suite 7 Rocklin, CA 95677

Telephone (916) 632-6800

Fax (916) 632-6812

October 26, 2007

Ms. Donna O'Brien Claims Representative State Compensation Insurance Fund P.O. Box 659011 2450 Venture Oaks Way, Suite 500 Sacramento, CA 95833-3291

Re:

State Board of Equalization Mold Evaluation on 3rd, 11th, and 21st Floors; 450 N Street,

Sacramento, CA

Dear Ms. O'Brien:

This report presents results of the very limited mold investigation by Entek Consulting Group, Inc. (Entek) at the State of California Board of Equalization (BOE) located at 450 N Street in Sacramento, CA. Ms. Judy Knight, Return to Work Coordinator of BOE requested our services to collect air samples on the 3rd, 11th, and the 21st floors of the building following work by others on the 23rd floor of the building at the perimeter walls. The investigation by Entek was very limited in scope and included limited sampling on these three floors to assist you with evaluation of mold spore levels.

The onsite inspection by Entek was conducted on September 29, 2007. I met with Ms. Judy Knight who provided me access to all office spaces for sampling and who was present during all of the sample collection. Environmental sampling was conducted at the building which included collection and analysis of 16 air samples for total non-culturable mold spores, two settled dust samples collected from the carpet and evaluated for mold spores, one vacuum bulk sample collected from the surface of a portable Honeywell HEPA filtered air cleaner for evaluation of particle identification by direct microscopic examination, and three bulk samples of suspect mold for evaluation of mold growth by direct microscopic examination.

Included in the total number of air samples indicated above, there were three air samples collected for nonculturable mold spores outside the building for comparison to the interior samples. The following is a discussion of each sampling technique and the results of the findings.

Air Sampling Results

Non-Culturable Mold Spores

Air sampling was conducted to evaluate non-culturable mold spores only on three floors in the building and was accomplished by collecting air samples onto "Air-O-Cell" sampling cassettes. The air sample is collected onto a coated plastic strip and visually evaluated by the analyst for all spores which stick to the coated slide. Since this technique includes evaluation for both non-culturable and culturable spores, the results will generally be higher than the sampling technique for culturable mold spores using the Anderson N6 impaction sampler, which relies on growth of spores onto a media.

There were 16 air samples collected and analyzed for non-culturable mold spores on this investigation, which included 13 air samples inside the building, and three air samples outside the building at the north side of the building by the picnic tables for comparison to the air samples collected inside. All of the sample times are noted on the chain of custody form for each location.

Asbestos Mold INDOOR AIR QUALITY L.FAD Noise Monitoring **TRAINING** HEALTH AND SAFETY AUDITS



Ms. Donna O'Brien State Compensation Insurance Fund October 26, 2007 Page Two

The samples were assigned a unique sample number, and sent to Environmental Microbiology Laboratory (EML) in Sacramento, CA, where they were evaluated by an analyst. The total concentration of mold spores inside the building ranged between 27 spores/M³ and 114 spores/M³, averaging 61 spores/M³. The average spore concentration on each of the different floors of the building tested are as follows: 3rd Floor --- 68 spores/M³, 11th Floor --- 35 spores/M³, and the 21st Floor --- 94 spores/M³. For comparison to the interior samples, the three outside air sample concentration of total mold spores were 4,850 spores/M³, 5,437 spores/M³, and 6,564 spores/M³, averaging 5,617 spores/M³.

The concentration of mold spores inside of the building at the areas tested compared to the concentration of mold spores outside of the building was significantly lower ranging between 0.5% to 2% of the outside levels.

It is also important to evaluate the distribution of mold spores seen inside a building compared to the outside air. If there is a significant increase in one or more individual spore types seen inside a building compared to the outside flora, it may be indicative of a mold source inside the building. The primary mold genera found in the outside air samples were *Cladosporium* followed by *Penicillium/Aspergillus* type spores, and Basidiospores (comprised primarily of mushroom type spores). Inside of the building, *Cladosporium* was also found to be the predominant genera detected; however, at concentrations much lower than that detected outdoors. There were no concentrations of specific genera of mold spores detected inside of the areas tested that were elevated above outside mold spore levels. This is also supportive of no significant mold source in the areas tested that would be contributing excessive spores to the office spaces.

MoldRANGE™ Extended Outdoor Comparison Report and MoldSTAT™ Supplementary Statistical Spore Trap Reports

Attached to each set of laboratory reports is additional information provided by EML regarding mold spore concentrations typically found outdoors during the month sampled for comparison to the results from our testing. The MoldRANGE™ Extended Outdoor Comparison report provides a review of a large data base of air samples collated by EML for locations across the United States for comparison to air sampling on any given day. This large data base of MoldRANGE™ provides a secondary comparison to the air samples collected by Entek for greater assurance of the types of mold spores expected and actually detected on the air samples. The results of the air samples on our investigation found concentrations of similar mold spore types as expected and seen in other areas of the country.

Also provided by EML are the MoldSTAT™ Supplementary Statistical Spore Trap Reports which compare each of the indoor air samples to the outside air samples collected on the day of sampling. This statistical evaluation provides a review of the comparison of the total mold spore concentration and type of mold spores detected inside the building to that detected outside the building. The "Mold Score" analysis provided by EML in their reports provides a relative "score" ranging between 100 and 300 using a statistical algorithm method developed by EML. A "score" of 100 is considered low and indicates or supports the premise that the concentration and types of mold spores detected on the air sample has a greater likelihood of coming from outside of the building, from an outside source. A score of 300 is considered high and indicates a greater likelihood of the mold spores originating from inside of the building. All of the "Mold Scores" on the air samples inside of the building were scored "low", which is indicative of no significant mold source inside of the spaces tested that would be contributing excessive mold spores into the occupied spaces.

There will always be some variability found in air sampling, hour by hour, day by day, month by month, and especially during different seasons. There will also be variability in sampling due to the randomness in distribution of spores in the air, doors and windows being open, and intake and filtration by the heating, ventilating, and air-conditioning (HVAC) system. The results of the air sampling on this investigation demonstrate this wide range in variability in mold spores measured both inside and outside of the building.



Ms. Donna O'Brien State Compensation Insurance Fund October 26, 2007 Page Three

It is normal and typical to find the presence or absence of a few genera in small numbers with this type of sampling. To illustrate the wide range in mold spore levels sometimes found, the outside air concentrations of mold spores detected on May 17, 2007 averaged 1,428 spores/M³, compared to the average outside mold spore concentration of 5,617 spores/M³ detected on September 29, 2007.

On the other hand, the interior mold spore levels inside of the building did not vary greatly with average mold spore concentration on May 17, 2007 at 110 spores/M³, compared to the average indoor mold spore concentration of 61 spores/M³ detected on September 29, 2007.

The mechanical HVAC fan unit for the three floors tested was operating during the air sampling for non-culturable mold spores. Prior to our sampling, we met with one of the engineering staff in the elevator and he assured that the air was operating on the three floors we were testing. This was also verified by observing air leaving the supply registers located at the ceiling systems in the office spaces. The air sampling was conducted on a Saturday, so there were only a handful of employees we happened to see inside of the building during my visit.

Other Biological Particles Results by Non-viable Methodology

Also evaluated on the air samples collected onto the Air-O-Cell cassettes were other airborne particles from other sources including pollen, plant, animal (primarily skin cells), fungi, and other non-biological particles such as glass fiber, soot, starch and synthetic fibers. The primary airborne particles identified in all of the indoor air samples were epithelial skin cells, due to normal shedding of skin cells by occupants.

Other particles detected in lessor amounts inside of the building were glass fibers, cotton fibers (from clothes), and trichomes (plant hairs), which may be due to some of the plants found inside of the various offices.

Overall, the total biological particulate loading found on the 13 air samples inside of the building on the three floors were low, which is reflective most likely of the quality filtration associated with the mechanical heating, ventilating, and air-conditioning system.

Bulk Sample Results

Carpet Dust Samples

At the request of Ms. Judy Knight, two samples of carpet dust were collected to evaluate mold spore loading in the carpet in the Law Library located on the 22nd floor and in the open office area in Room 2206 near cubicle #095. The purpose of this limited sampling was to compare the total mold spore loading in these two locations following the carpet cleaning that was conducted since my initial sampling on May 17, 2007 in the same locations.

Sampling of the dust was performed using a 0.45 micron mixed cellulose ester filter attached to a high volume pump with tygon tubing at a flow rate of greater than 15 liters per minute. The plastic top section of the filter cassette was removed, and the open filter cassette was placed onto the carpet surfaces being vacuumed. The sample was collected as a composite sample comprised of at least four different approximately one square foot locations until a large enough bulk sample was collected into the cassette, similar to the method used on May 17, 2007.



Ms. Donna O'Brien State Compensation Insurance Fund October 26, 2007 Page Four

The samples were individually labeled and submitted to EML of San Bruno, CA for evaluation by a staff analyst. The samples were diluted in a solution, and each were plated onto three different petri dishes containing different media, which included Cellulose, Malt Extract Agar (MEA), and Dichloran glycerol (DG 18). After several days of growth, the analyst identified the mold genera and species (if possible) and concentration. The following table provides a summary of the results of the carpet dust sampling conducted on May 17, 2007 and the more recent sampling on September 29, 2007 following cleaning of the carpets.

Table of Sample Results of Carpet Dust Analysis for Mold Spores on May 17, 2007 Before Carpet was Cleaned and on September 29, 2007 After Carpet was Cleaned

Location	Results of May 17, 2007 Prior to Carpet Cleaning (cfu/gram)	Results of September 29, 2007 After Carpet Cleaning (cfu/gram)
Room 2217 Law Library	7,232,400	246,000
Room 2206 near Cubicle #095	112,000	12,800

There are no standards for mold spore levels in carpet dust; however, they can assist in evaluating settled particles which may have been deposited over many weeks, months, or years depending upon the frequency and thoroughness of the carpet cleaning. Some of the mold spores considered primary water indicators include *Aspergillus*, *Penicillium*, *Chaetomium*, *Stachybotrys*, *Fusarium*, and *Ulocladium* to name a few of the more common mold genera. There were very few *Aspergillus*, *Penicillium* and *Fusarium* colonies observed and no *Stachybotrys*, *Ulocladium*, or *Chaetomium* colonies observed. It is common to find low levels of various mold genera with this type of sampling.

The results of these two carpet dust samples collected in September suggest the carpet cleaning conducted following the initial sampling in May have resulted in a significant reduction in total mold spore loading. Although only two samples were collected, it is indicative of the importance of good cleaning techniques described in my report of June 15, 2007. Assuming the carpet has been cleaned in other office areas in a similar fashion as in the two areas tested on September 29th, if additional carpet dust samples were collected, I would suspect similar results reflecting a similar reduction in the total mold spore burden.

Results of Bulk Samples of Suspect Mold

Three bulk samples were collected of suspect mold from building surfaces; two from the 22nd floor and one from the 23rd floor of the building. Samples were collected using clear scotch tape pressed onto the suspect discoloration to lift off the suspect mold and placed onto a clear microscope slide. Samples were identified with a unique sample number, placed in a sealed zip lock bag and submitted to EML of Sacramento, CA for evaluation by an analyst.

One sample was collected from the base of the south gypsum wall near the south east corner across from cubicle #095. Stachybotrys and Acremonium mold was observed by the laboratory. A second sample was collected from the base of the west gypsum wall near the door of Room 2238 and Penicillium/Aspergillus type spores were observed. The third sample was collected from the south perimeter wall east of cubicle #33 on the 23rd floor and Aspergillus type mold was observed by the laboratory. These three samples were collected in locations identified as having a mold issue by other investigators prior to my onsite visit. The one sample collected at the lower perimeter wall on the 23rd floor was able to be collected, since the office furniture had been removed from the perimeter wall prior to my visit allowing access.



Ms. Donna O'Brien State Compensation Insurance Fund October 26, 2007 Page Five

Results of Particulate Evaluation Associated with a Honeywell HEPA Filter Unit

During my visit, Ms. Judy Knight requested an evaluation of the particulate associated with the surface of a portable Honeywell HEPA filter unit that was taken out of service and placed at Judy's cubicle.

Apparently, one employee was concerned with the particulate matter on the outside of the plastic housing associated with this portable filtration device. I collected one sample of settled particulate from the exterior surface of the Honeywell unit onto a 0.45 micron mixed cellulose ester (MCE) filter housed in a plastic cassette with a small piece of Tygon tubing attached to the end of the cassette. The micro-vacuum bulk sample was collected using a high volume sample pump to act as a vacuum cleaner to collect the sample. The sample was submitted to Forensic Analytical Specialties, Inc. of Hayward, CA for particle identification using polarized light microscopy (PLM).

Attached are the analytical results of the particle analysis, which includes a breakdown of the *Fibrous* and *Non-fibrous* fractions in the sample. Of the *fibrous* fraction, the sample was found to have major amounts (greater than 10%) of cotton fibers and cellulose. The primary source of cotton is from clothing worn by the employees. There were minor amounts (1-10%) of paper dust found on the sample. There were trace amounts (< 1%) of synthetic fibers, wool, mineral wool, trichomes (plant hairs), and cat hair.

Of the *non-fibrous* fraction, there were major amounts of epithelial (skin) cells detected, which is very common inside of occupied buildings. There were minor amounts of clear isotropics, which may be comprised of glass chips or aluminum corrosion products. There were trace amounts of pollen, iron oxide, limestone, feldspars, opaques, organic debris, paint chips, quartz, spray paint, starch, salt, and phenolic foam.

These analytical findings are very typical of indoor air particulate found on many other investigations by Entek and they will vary somewhat in the composition percentages in different buildings, but generally the variety and distribution of the different biological, mineral, and man-made particles is common. The particulate on the surface of the Honeywell fan unit is reflective of particles brought in from the outside environment or generated inside of the building, which are eventually released from the supply ducts during operation of the fan system due to air flow and vibration or brought in by the occupants (such as cat hair for example).

Summary

The air sampling for mold spores by Entek Consulting Group, Inc. was limited in scope and included the 3rd, 11th, and the 21st floors of the building following work by others on the 23rd floor at the perimeter walls. Air sampling for total mold spores by non-culturable methods found levels to be much lower than the air samples collected outside of the building, indicative of no major mold source inside the areas tested that might be contributing significant mold spores into the occupied spaces.

There were 13 air samples collected inside of the building on this limited survey. The total concentration of mold spores inside the areas tested ranged between 27 spores/M³ and 114 spores/M³, averaging 61 spores/M³. For comparison, the average concentration of total mold spores detected inside of the areas tested during my investigation on May 17, 2007 was 110 spores/m³. The results of the testing on September 29, 2007 are slightly lower, but within the same order of magnitude as found in May. The lower results may be due to a combination of the carpet cleaning efforts and the lack of employees present during the sampling in September, since sampling was conducted on a Saturday when very few employees were present with less activity by occupants such as walking around.

There were low levels of other biological particulate observed on the 13 air samples collected inside of the areas tested. The type of biological particulate is typical of indoor environments and will vary greatly with the type of mechanical HVAC equipment and filtration present. The more sophisticated the filtration associated with the HVAC system, the lower the levels of total particulate including biological particulate. The low



Ms. Donna O'Brien State Compensation Insurance Fund October 26, 2007 Page Six

concentrations of other biological particulate and mold spores illustrate the high efficiency in the filtration associated with the HVAC system at the Board of Equalization.

The bulk samples collected of suspect mold on several building components confirmed the presence of mold growth. Samples were collected from the base of gypsum wallboard in two samples and one sample was collected approximately 12-15" above floor level at the perimeter south wall on the 23rd floor where there has been a history of water intrusion.

At the time of my investigation, the 23rd floor was under various degrees of construction with plastic sheeting at the hallways to restrict access to the construction areas. It is my understanding painting of the walls was one of the objectives on the 23rd floor, resulting in the findings of mold after office furniture was removed from the perimeter walls and after removal of one or more layers of the gypsum wallboard (sheet rock). No air sampling was conducted on the 23rd floor, since there were open wall systems observed and the areas were cordoned off with the plastic barriers.

The results of the two carpet dust sample analysis found a significant reduction in total mold spores compared to the testing conducted on May 17, 2007. The purpose of the carpet dust samples was to evaluate the total mold spore loading following the carpet cleaning. The total concentration of mold spores in the sample collected at the Law Library on September 29, 2007 were less than 1% of the total mold spore loading found in our sampling on May 17, 2007. Similarly, the total concentration of mold spores in the sample collected in Room 2206 near cubicle #095 on September 29, 2007 were less than 5% of the total mold spore loading found in our sampling on May 17, 2007.

These reductions in total mold spore loading in the carpet illustrate the significance of thorough cleaning of carpeted areas inside of buildings. The frequency of cleaning, thoroughness of cleaning, and cleaning techniques (HEPA vacuuming or steam extraction cleaning) is critical in reducing total particulate burden (including mold spores) in carpets. Continual efforts should be in place to address the carpet cleaning in the building on all floors, not just the various floors of recent concern.

The results of the analysis of the settled dust associated with the Honeywell HEPA filter cleaning unit is reflective of normal particulate seen inside of buildings. The plastic housing on many office equipment can also become electrostatically charged, further attracting particulate onto the surfaces. Settled brown or black particulate seen on supply or return registers, computer terminals, and other office equipment typically will have a similar profile of various particles as seen in the limited sampling conducted on September 29, 2007.

It has been my pleasure working with you on this investigation. Thank you for choosing Entek Consulting Group, Inc. for your environmental needs. Please call me at ((16) 632-6800 if you have any questions regarding this report.

Sincerely,

Richard Beall, CIH, CSP

President

Enclosures

C:\Entek\Clients\StateComplnsuranceFund\07-534A Board of Equalization Mold\Mold Investigation Report.wpd



NON-CULTURABLE MOLD SPORE

AIR SAMPLING RESULTS



ENTEK CONSULTING GROUP, INC.

4200 Rocklin Road, Suite 7 Rocklin, CA 95677 (916) 632-6800 (916) 632-6812 FAX

Page 1 of 2

TABLE OF AIR SAMPLING RESULTS NON-CULTURABLE MOLD SPORES AND OTHER BIOLOGICAL PARTICULATE

Date of Sampling: 09/29/07

Lab: EML - Sacramento

Job Number: 07-534A

Turnaround Time:

Standard

ENTEK

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization 450 N Street Sacramento, CA

SAMPLE NUMBER	SAMPLE LOCATION	TIME ON/OFF	RESULTS SPORES/M³
ECG-07-534A-500	3 rd Floor Room 322 Cubicle #56	13:52:52 14:02:52	94
ECG-07-534A-501	3 rd Floor Room 317 Cubicle #35 at Isle Way	13:48:38 13:58:38	102
ECG-07-534A-502	3 rd Floor Room 311 Staging Area #129	14:12:12 14:22:12	34
ECG-07-534A-503	3 rd Floor Room 327 Cubicle #94	14:09:45 14:19:45	41
ECG-07-534A-504	11 th Floor Room 1104 Cubicle #56	14:29:48 14:39:48	27
ECG-07-534A-505	11 th Floor Room 1104 Cubicle #32	14:33:53 14:43:53	60
ECG-07-534A-506	11 th Floor Room 1104 Cubicle #28	14:48:44 14:58:44	27
ECG-07-534A-507	11 th Floor Room 1104 Cubicle #4.0	14:51:27 15:01:27	27
ECG-07-534A-508	21 st Floor Room South East Area Cubicle #3	15:43:53 15:53:53	107
ECG-07-534A-509	21 st Floor Room South West Area Cubicle #30	15:29:45 15:39:45	114
ECG-07-534A-510	21 st Floor Room North East Area Cubicle #119	15:09:19 15:19:19	34
ECG-07-534A-511	21 st Floor Room West Area Cubicle #55	15:27:05 15:37:05	87
ECG-07-534A-512	21 st Floor Room North Area Cubicle #109	15:11:37 15:21:37	34



ENTEK CONSULTING GROUP, INC. 4200 Rocklin Road, Suite 7 Rocklin, CA 95677 (916) 632-6800 (916) 632-6812 FAX

Page 2 of 2

TABLE OF AIR SAMPLING RESULTS NON-CULTURABLE MOLD SPORES AND OTHER BIOLOGICAL PARTICULATE

Date of Sampling: 09/29/07

Lab:

EML - Sacramento

Job Number: 07-534A

Turnaround Time:

Standard

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization

450 N Street Sacramento, CA

SAMPLE NUMBER	SAMPLE LOCATION	TIME ON/OFF	RESULTS SPORES/M³
ECG-07-534A-513	Outside Ambient Air - North Side Outside of Cafeteria	13:30:47 13:40:47	5,437
ECG-07-534A-514	Outside Ambient Air - North Side Outside of Cafeteria	16:03:00 16:13:00	4,850
ECG-07-534A-515	Outside Ambient Air - North Side Outside of Cafeteria	16:03:00 16:13:00	6,564

Sample flow rates for all air sampling at 15.0 liters per minute calibrated using a rotameter that was calibrated by a primary standard (BIOS, DryCal, DC-Lite, Serial Number 3518, Model Number DCL-H Rev. 1.08). Total volume collected was 150 liters of air.

All indoor samples were collected onto Zefon "Air-O-Cell" cassettes for a total of 10 minutes each.

Outside Ambient Air Conditions (Temperature & Wind):

°F; Winds, Clear Skies

HVAC fan on during air sampling.

Z:\Clients\State Comp Ins Fund\07-534A Board of Equal 450 N Street\NonVia.Tbl.500



Ratio of Indoor to Outdoor Mold Spore Concentrations Other Biological Particulate

Date of Sampling: 09/29/07

Lab:

EML - Sacramento

Job Number: 07-534A

Turnaround Time:

Standard

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization 450 N Street

Sacramento, CA

Location	Co	Mold Spore	sured
Room	CFU/M³	<u>[Inside]</u> [Outside]	% of [Outside]
Outside Ambient Air	5,617		
3 rd Floor Room 322 Cubicle #56	94	0.017	1.7
3 rd Floor Room 317 Cubicle #35 at Isle Way	102	0.018	1.8
3 rd Floor Room 311 Staging Area #129	34	0.006	0.6
3 rd Floor Room 327 Cubicle #94	41	0.007	0.7
11 th Floor Room 1104 Cubicle #56	27	0.005	0.5
11 th Floor Room 1104 Cubicle #32	60	0.011	1.1
11 th Floor Room 1104 Cubicle #28	27	0.005	0.5
11 th Floor Room 1104 Cubicle #4.0	27	0.005	0.5
21 st Floor Room South East Area Cubicle #3	107	0.019	1.9
21 st Floor Room South West Area Cubicle #30	114	0.020	2.0
21 st Floor Room North East Area Cubicle #119	34	0.006	0.6
21st Floor Room West Area Cubicle #55	87	0.015	1.5
21 st Floor Room North Area Cubicle #109	34	0.006	0.6

Z:\Clients\State Comp Ins Fund\07-534A Board of Equal 450 N Street\MoldRatio.500.wpd

^{*}Outside Ambient Air is a average of samples ECG-07-534A-0513, ECG-07-534A-0514, & ECG-07-534A-0515. Total of samples were 16,851 \div 3 making 5,617 the overall average.

ENTEK CONSULTING GROUP, INC.

4200 Rocklin Road, Suite 7 Rocklin, CA 95677 (916) 632-6800 (916) 632-6812 FAX

TABLE OF AIR SAMPLING RESULTS NON-CULTURABLE MOLD SPORES AND OTHER BIOLOGICAL PARTICULATE

Date of Sampling: 09/29/07

Lab:

EML - Sacramento

Job Number: 07-534A

Turnaround Time:

Standard

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization

450 N Street Sacramento, CA

SAMPLE NUMBER SAMPLE LOCATION TIME RESULTS ON/OFF SPORES/M3 3rd Floor Room 322 Cubicle #56 ECG-07-534A-500 13:52:52 14:02:52 ECG-07-534A-501 3rd Floor Room 317 Cubicle #35 at Isle Way 13:48:38 13:58:38 ECG-07-534A-502 3rd Floor Room 311 Staging Area #129 14:12:12 14:22:12 ECG-07-534A-503 3rd Floor Room 327 Cubicle #94 14:09:45 14:19:45 ECG-07-534A-504 11th Floor Room 1104 Cubicle #56 14:29:48 14:39:48 11th Floor Room 1104 Cubicle #32 ECG-07-534A-505 14:33:53 14:43:53 ECG-07-534A-506 11th Floor Room 1104 Cubicle #28 14:48:44 14:58:44 ECG-07-534A-507 11th Floor Room 1104 Cubicle #4.0 14:51:27 15:01:27 ECG-07-534A-508 21st Floor Room South East Area Cubicle #3 15:43:53 15:53:53 ECG-07-534A-509 21st Floor Room South West Area Cubicle #30 15:29:45 15:39:45 ECG-07-534A-510 21st Floor Room North East Area Cubicle #119 15:09:19 15:19:19 ECG-07-534A-511 21st Floor Room West Area Cubicle #55 15:27:05 15:37:05 ECG-07-534A-512 21st Floor Room North Area Cubicle #109 15:11:37 15:21:37

(916) 632-6800 (916) 632-6812 FAX

TABLE OF AIR SAMPLING RESULTS NON-CULTURABLE MOLD SPORES AND OTHER BIOLOGICAL PARTICULATE

Date of Sampling: 09/29/07

Lab:

EML - Sacramento

Job Number: 07-534A

Turnaround Time:

Standard

Page 2 of 2

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization

450 N Street Sacramento, CA

SAMPLE LOCATION	TIME	RESULTS
	T ON/OFF	SPORES/M ³
Outside Ambient Air - North Side Outside of Cafeteria	13:30:47 13:40:47	
Outside Ambient Air - North Side Outside of Cafeteria	16:03:00	-
Outside Ambient Air - North Side Outside of Cafeteria	16:03:00	
	Outside Ambient Air - North Side Outside of Cafeteria Outside Ambient Air - North Side Outside of Cafeteria Outside Ambient Air - North Side Outside of	Outside Ambient Air - North Side Outside of Cafeteria 13:30:47 Outside Ambient Air - North Side Outside of 16:03:00 Cafeteria 16:03:00 Cafeteria 16:03:00 Cutside Ambient Air - North Side Outside of 16:03:00

Sample flow rates for all air sampling at 15.0 liters per minute calibrated using a rotameter that was calibrated by a primary standard (BIOS, DryCal, DC-Lite, Serial Number 3518, Model Number DCL-H Rev. 1.08). Total volume collected was 150 liters of air.

All indoor samples were collected onto Zefon "Air-O-Cell" cassettes for a total of 10 minutes each.

Outside Ambient Air Conditions (Temperature & Wind):

°F; Winds, Clear Skies

HVAC fan on during air sampling.

Z:\Clients\State Comp Ins Fund\07-534A Board of Equal 450 N Street\NonVia.Tbl.500

Delivered by:	Meren	Date: 12:05 PM Time: 10-1-07
Received by:	Siehon	Date: 12 05 / Time: 10 / / 6

Client: Entek Consulting Group C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:		-534A-500: r room 322	ECG-07	-534A-501: r room 317	ECG-07	-534A-502:	ECG-07	-534A-503
	cubicle #56		cubicle	#35 at Isle Way	staging	r room 311 area #129		r room 327 cle #94
Comments (see below)	1	Vone	1	None		Vone	None	
Lab ID-Version‡:	149	9735-1	149	9736-1	149	9737-1	149	9738-1
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria							1	7
Arthrinium								
Ascospores*			1	. 27				
Aureobasidium								
Basidiospores*							-	
Bipolaris/Drechslera group								
Chaetomium					· ·			
Cladosporium	2	53	1	27	1	27	1	27
Curvularia			1	7		21	т	21
Epicoccum				, , , , , , , , , , , , , , , , , , , ,				
Fusarium								
Nigrospora	1	7						
Oidium								
Other brown								
Penicillium/Aspergillus types†	1	27	1	27			***	
Pithomyces		,						
Rusts*					1	7		
Smuts*, Periconia, Myxomycetes*	1	7	1	7	.		1	7
Stemphylium								
Torula								*****
Ulocladium			1	7				
Unknown				,				
Background debris (1-4+)††	2+		2+		2+		2+	· · · · · · · · · · · · · · · · · · ·
Hyphal fragments/m3	13		7		13		7	
Pollen/m3	13		13		13		< 7	
Skin cells (1-4+)	1+		1+		1+		1+	
Sample volume (liters)	150		150		150		150	
TOTAL SPORE/m3		94		102		34	130	41

Comments:

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi.

^{*}Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sportnating rungi. Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.
† The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††} Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be actually higher than reported. Background debris also affects the reporting limit for some spore types. The reporting limit is dependent on spore size, background debris, sample volume, and the percentage of the trace analyzed. It is important to account for sample volumes when evaluating dust levels.

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	11th f	-534A-504: loor room ubicle #56	11th f	7-534A-505: loor room ubicle #32	11th f	7-534A-506: loor room ubicle #28	11th f	7-534A-507: loor room ubicle #4.0
Comments (see below)	1	Vone	1	Vone	1	Vone	None	
Lab ID-Version‡:	149	9739-1	149	1499740-1		9741-1	1499742-1	
			spores/m3					
Alternaria	2000000	550100/1115	1	7	raw ct.	spores/m3	raw ct.	spores/m3
Arthrinium								
Ascospores*								ļ
Aureobasidium								
Basidiospores*								
Bipolaris/Drechslera group								
Chaetomium								
Cladosporium	1	27	2	53	1	27	1	27
Curvularia				33		21		27
Epicoccum								
Fusarium								
Nigrospora								
Oidium								
Other brown								
Penicillium/Aspergillus types†								
Pithomyces								
Rusts*								
Smuts*, Periconia, Myxomycetes*								
Stachybotrys								
Stemphylium								
Torula								
Ulocladium								
Unknown								
Background debris (1-4+)††	2+		2+		2+		1+	
Hyphal fragments/m3	7		7		7		< 7	
Pollen/m3	7		< 7		< 7		< 7	
Skin cells (1-4+)	1+		1+		1+		< 1+	
Sample volume (liters)	150		150		150		150	
FOTAL SPORE/m3 Comments:		27		60	150	27	120	27

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi. Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and

^{††} Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be actually higher than reported. Background debris also affects the reporting limit for some spore types. The reporting limit is dependent on spore size, background debris, sample volume, and the percentage of the trace analyzed. It is important to account for sample volumes when

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	ECG-07-534A-508: 21st floor room south east area cubicle #3		ECG-07 21st fl south	7-534A-509: oor room west area icle #30	21st fl north	7-534A-510: loor room east area cle #119	ECG-07-534A-511: 21st floor room west area cubicle # 55		
Comments (see below)	1	Vone	1	Vone	1	Vone	None		
Lab ID-Version‡:	149	9743-1	149	9744-1	149	9745-1		9746-1	
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	raw ct	spores/m?	
Alternaria			1	7			1417 00.	3p0103/111.	
Arthrinium									
Ascospores*	11	27							
Aureobasidium									
Basidiospores*					-		· · · · · · · · · · · · · · · · · · ·		
Bipolaris/Drechslera group				-					
Chaetomium							1	7	
Cladosporium	2	53	4	107	1	27	3		
Curvularia			••••••	107	<u> </u>	21		80	
Epicoccum									
Fusarium					-				
Nigrospora									
Oidium									
Other brown						-			
Penicillium/Aspergillus types†	1	27							
Pithomyces		27			<u> </u>		-		
Rusts*		-							
Smuts*, Periconia, Myxomycetes*		-			1				
Stemphylium						7			
Γorula									
Ulocladium									
Unknown									
Background debris (1-4+)††	2+		2+						
Hyphal fragments/m3	< 7		< 7		2+		1+		
Pollen/m3	< 7		< 7		< 7		< 7		
Skin cells (1-4+)	1+		1+		< 7		7		
Sample volume (liters)	150		150		1+		1+		
TOTAL SPORE/m3		107	130	114	150		150		
Comments:		10/		114		34		87	

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi.

Most of these spore types are not seen with culturable memous (Andersen sampling), annough some may appear as non-sportnating rung.

Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.

† The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and

^{††} Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be actually higher than reported. Background debris also affects the reporting limit for some spore types. The reporting limit is dependent on spore size, background debris, sample volume, and the percentage of the trace analyzed. It is important to account for sample volumes when

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of

[‡] A "Version" greater than 1 indicates amended data.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	21st fl	-534A-512: oor room ea cubicle #	Outside north sid	ambient air, e outside of	Outside north sid	7-534A-514: ambient air, le outside of	Outside north sid	7-534A-515 ambient air le outside o	
	109		Ca	feteria	ca	feteria	cafeteria		
Comments (see below)	1	Vone	<u> </u>	None	<u> </u>	None			
Lab ID-Version‡:	149	9747-1		9748-1		9749-1	140	A 9750-1	
	raw ct.	spores/m3		spores/m3					
Alternaria		БРОГОВ/ПІЗ	9	60		spores/m3	raw ct.	spores/m.	
Ascospores*			3	80	6	40	7	47	
Aureobasidium				- 00	2	53	3	80	
Basidiospores*			14	373	10	267			
Bipolaris/Drechslera group				3/3	10	267	9	240	
Chaetomium					1	7			
Cladosporium	1	27	175	4,670	$\frac{1}{160}$		1774	4.640	
Curvularia	-		1	7	3	4,270	174	4,640	
Epicoccum			2	13	2	20 13	1		
Fusarium						13	1	7	
Nigrospora			3	20	14	93			
Oidium					1 1	7	7	47	
Other brown			1	7	1	7	2	13	
Penicillium/Aspergillus types†			4	107			104		
Pithomyces				107			194	1,430	
Rusts*			4	27	2	12			
Smuts*, Periconia, Myxomycetes*	1	7	11	73	6	13	4	27	
Stemphylium					_0	40	3	20	
Torula									
Ulocladium					3	20			
Unknown					_3	20	2	13	
Background debris (1-4+)††	1+		3+		3+				
Hyphal fragments/m3	< 7		73		53		3+		
Pollen/m3	< 7		67		40		13 13		
Skin cells (1-4+)	1+		1+		1+		13		
Sample volume (liters)	150		150		150		150		
TOTAL SPORE/m3 Comments: A) 187 of the raw count Pen		34		5,437		4,850	130	6,564	

Comments: A) 187 of the raw count Penicillium/Aspergillus type spores were present as a single clump.

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi. Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.

† The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing

characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††} Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be actually higher than reported. Background debris also affects the reporting limit for some spore types. The reporting limit is dependent on spore size, background debris, sample volume, and the percentage of the trace analyzed. It is important to account for sample volumes when

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of

[‡] A "Version" greater than 1 indicates amended data.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldRANGE™: Extended Outdoor Comparison

Outdoor Location: ECG-07-534A-513, Outside ambient air, north side outside of cafeteria

Fungi Identified	Outdoor	Typic	al Outdo	or Data by				Data by I	ocation	
	data		Month:	September		State: CA				
	spores/m3	low	med	high	freq %	low	med	high	freq %	
Generally able to grow indoors*									1104 70	
Alternaria	60	7	38	540	66	7	27	230	61	
Bipolaris/Drechslera group	· _	7	13	210	26	7	13	120		
Chaetomium	-	7	13	120	17	7	13		14	
Cladosporium	4,670	53	800	12,000	98	53	640	110	19	
Curvularia	7	7	27	820	31	7	13	6,500 190	98	
Epicoccum	13	7	22	430	28	7	13	160	7	
Nigrospora	20	7	20	360	25	7	13	180	21	
Other brown	7	7	13	110	39	7	13	180 88	8	
Penicillium/Aspergillus types	107	47	290	3,700	92	50	210		38	
Stachybotrys	-	7	13	280	4	7	13	2,600 350	89	
Torula	_	7	13	120	16	7	13	150	5	
Ulocladium	-	7	13	160	9	7	13		13	
Seldom found growing indoors**				100		,	13	110	9	
Ascospores	80	13	170	5,500	80	13	110	1 700	72	
Basidiospores	373	20	390	23,000	96	13	270	1,700	73	
Oidium	_	7	13	170	15	7		7,000	95	
Rusts	27	7	22	340	29	7	13 13	200	20	
Smuts, Periconia, Myxomycetes	73	10	53	730	81	8	40	270	29	
TOTAL SPORES/M3	5,437			750	01		40	480	72	

† The Typical Outdoor Data by Date represents the typical outdoor spore levels across North America for the month indicated. The last column represents the frequency of occurrence. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 2.5% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

‡ The Typical Outdoor Data by Location represents the typical outdoor spore levels for the region indicated for the entire year. As with the Typical Outdoor Data by Date, the four columns represent the frequency of occurrence and the typical low, medium, and high concentration values for the spore type indicated. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

*The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

**These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldRANGETM: Extended Outdoor Comparison

Outdoor Location: ECG-07-534A-514, Outside ambient air, north side outside of cafeteria

Fungi Identified	Outdoor	Typic	al Outdo	or Data by	Date†	Typical	Outdoor	Data by I	ocation t	
	data		Month:	September		State: CA				
	spores/m3	low	med	high	freq %	low	med	high	freq %	
Generally able to grow indoors*									1 111	
Alternaria	40	7	38	540	66	7	27	230	61	
Bipolaris/Drechslera group	-	7	13	210	26	7	13	120	14	
Chaetomium	7	7	13	120	17	7	13	110	19	
Cladosporium	4,270	53	800	12,000	98	53	640	6,500	98	
Curvularia	20	7	27	820	31	7	13	190	7	
Epicoccum	13	7	22	430	28	7	13	160	21	
Nigrospora	93	7	20	360	25	7	13	180	8	
Other brown	7	7	13	110	39	7	13	88	38	
Penicillium/Aspergillus types	-	47	290	3,700	92	50	210	2,600	89	
Stachybotrys	-	7	13	280	4	7	13	350	5	
Torula	-	7	13	120	16	7	13	150	13	
Ulocladium	20	7	13	160	9	7	13	110	9	
Seldom found growing indoors**						•	13	110	9	
Ascospores	53	13	170	5,500	80	13	110	1,700	73	
Basidiospores	267	20	390	23,000	96	13	270	7,000	95	
Oidium	7	7	13	170	15	7	13	200	20	
Rusts	13	7	22	340	29	7	13	270	20 29	
Smuts, Periconia, Myxomycetes	40	10	53	730	81	8	40	480	29 72	
TOTAL SPORES/M3	4,850				- 01		70	400	12	

† The Typical Outdoor Data by Date represents the typical outdoor spore levels across North America for the month indicated. The last column represents the frequency of occurrence. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 2.5% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

‡ The Typical Outdoor Data by Location represents the typical outdoor spore levels for the region indicated for the entire year. As with the Typical Outdoor Data by Date, the four columns represent the frequency of occurrence and the typical low, medium, and high concentration values for the spore type indicated. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

*The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

**These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldRANGETM: Extended Outdoor Comparison

Outdoor Location: ECG-07-534A-515, Outside ambient air, north side outside of cafeteria

Fungi Identified	Outdoor	Typic	al Outdoo	or Data by	Date†	Typical	Outdoor	Data by L	ocation‡
	data	Month: September			State: CA				
	spores/m3	low	med	high	freq %	low	med	high	freq %
Generally able to grow indoors*									
Alternaria	47	7	38	540	66	7	27	230	61
Bipolaris/Drechslera group	-	7	13	210	26	7	13	120	14
Chaetomium	-	7	13	120	17	7	13	110	19
Cladosporium	4,640	53	800	12,000	98	53	640	6,500	98
Curvularia	-	7	27	820	31	7	13	190	7
Epicoccum	7	7	22	430	28	7	13	160	21
Nigrospora	47	7	20	360	25	7	13	180	8
Other brown	-	7	13	110	39	7	13	88	38
Penicillium/Aspergillus types	1,430	47	290	3,700	92	50	210	2,600	89
Stachybotrys	-	7	13	280	4	7	13	350	5
Torula	-	7	13	120	16	7	13	150	13
Ulocladium	13	7	13	160	9	7	13	110	9
Seldom found growing indoors**									
Ascospores	80	13	170	5,500	80	13	110	1,700	73
Basidiospores	240	20	390	23,000	96	13	270	7,000	95
Oidium	13	7	13	170	15	7	13	200	20
Rusts	27	7	22	340	29	7	13	270	29
Smuts, Periconia, Myxomycetes	20	10	53	730	81	8	40	480	72
TOTAL SPORES/M3	6,564								

[†] The Typical Outdoor Data by Date represents the typical outdoor spore levels across North America for the month indicated. The last column represents the frequency of occurrence. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 2.5% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

[‡] The Typical Outdoor Data by Location represents the typical outdoor spore levels for the region indicated for the entire year. As with the Typical Outdoor Data by Date, the four columns represent the frequency of occurrence and the typical low, medium, and high concentration values for the spore type indicated. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

^{*}The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**}These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: ECG-07-534A-513: Outside ambient air, north side outside of cafeteria

				Outdoor sample spores/m3 Typical outdoor ranges								
Species detected		Outdoor	sample s	pores/m:	3	Typical	outdo	or ranges	Freq.			
	<100	1K	10K	>100K		(Nor	th An	nerica)	%			
Alternaria					60	7 -	27	- 37Ó	55			
Ascospores					80	13 -	160	- 4,300	76			
Basidiospores					373	13 -	320	- 14,000	93			
Cladosporium					4,670	53 -	530	- 8,300	95			
Curvularia					7	7 -	22	- 670	16			
Epicoccum	195				13	7 -	13	- 280	23			
Nigrospora					20	7 -	13	- 230	13			
Other brown	677 887 887				7	7 -	13	- 93	36			
Penicillium/Aspergillus types					107	27 -	210	- 2,600	87			
Rusts					27	7 -	13	- 280	23			
Smuts, Periconia, Myxomycetes	G. H				73	7 -	40	- 760	70			
Total					5,437	•			'			

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: ECG-07-534A-500: 3rd floor room 322 cubicle #56

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ent ratio** r/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)			
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Resu	lt: 0.5333	dF: 11 Result: 0.5727 Critical value: 0.5273 Outside Similar: Yes	Score: 104 Result: Low			
Species 1	Species Detected		Spores/m3					
		<100	1K	10K	>100K			
	Cladosporium	ELECTION OF THE PROPERTY OF TH			53			
	Nigrospora				7			
	Penicillium/Aspergillus types				7 27			
Smuts, P	Smuts, Periconia, Myxomycetes				7			
	Total				94			

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-501: 3rd floor room 317 cubicle #35 at Isle Way

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: (0.5882	dF: 12 Result: 0.4283 Critical value: 0.4965 Outside Similar: No	Score: 107 Result: Low	
Species 3	Species Detected			Spores/m3		
		<100	1K	10K	>100K	
	Ascospores	77218 77184			27	
	Cladosporium	WARE CARE			27	
	Curvularia				7	
Penici	llium/Aspergillus types				27	
Smuts, Periconia, Myxomycetes						
	Ulocladium					
	Total				102	

Location: ECG-07-534A-502: 3rd floor room 311 staging area #129

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor) Result: 0.3077		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes			dF: 11 Result: 0.5023 Critical value: 0.5273 Outside Similar: No	Score: 100 Result: Low	
Species 1	Detected			Spores/m3		
		<100	1K	10K	>100K	
	Cladosporium				27	
	Rusts					
	Total				34	

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-503: 3rd floor room 327 cubicle #94

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.4286	dF: 11 Result: 0.5364 Critical value: 0.5273 Outside Similar: Yes	Score: 104 Result: Low	
Species 1	Detected		Spores/m3		
		<100 1K	10K	>100K	
	Alternaria			7	
Cladosporium				27	
Smuts, Periconia, Myxomycetes					
	Total	2015 2016 2016		41	

Location: ECG-07-534A-504: 11th floor room 1104 cubicle #56

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	·	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1667	dF: 11 Result: 0.6273 Critical value: 0.5273 Outside Similar: Yes	Score: 100 Result: Low
Species 1	Detected	<100 1K	Spores/m3	. 1007
	Cladosporium Total		10K	>100K

Location: ECG-07-534A-505: 11th floor room 1104 cubicle #32

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3077		dF: 11 Result: 0.5477 Critical value: 0.5273 Outside Similar: Yes	Score: 102 Result: Low	
Species 1	Detected			Spores/m3	Jan	
		<100	1K	10K	>100K	
	Alternaria				7	
	Cladosporium				53	
	Total					

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-506: 11th floor room 1104 cubicle #28

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		correl	nan rank ation*** /outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1667		dF: 11 Result: 0.6273 Critical value: 0.5273 Outside Similar: Yes		Score: 100 Result: Low	
Species	Detected	-100		Spo	res/m3	:	
	Cladosporium	<100	1K		10K	>100K	
	Total	ORGANA				27	

Location: ECG-07-534A-507: 11th floor room 1104 cubicle #4.0

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1667		dF: 11 Result: 0.6273 Critical value: 0.5273 Outside Similar: Yes	Score: 100 Result: Low
Species 1	Detected			Spores/m3	
-		<100	1K	10K	>100K
	Cladosporium				27
	<u>Total</u>				27

Location: ECG-07-534A-508: 21st floor room south east area cubicle #3

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.4286	dF: 11 Result: 0.7182 Critical value: 0.5273 Outside Similar: Yes	Score: 104 Result: Low	
Species 1	Detected		Spores/m3		
		<100 1K	10K	>100K	
	Ascospores	100 miles		27	
Cladosporium				53	
Penici	Penicillium/Aspergillus types				
	Total			107	

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTAT™: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-509: 21st floor room south west area cubicle #30

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)		
Result: 2%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3077		Resul Critical v	dF: 11 Result: 0.5477 Critical value: 0.5273 Outside Similar: Yes		re: 102 llt: Low	
Species 3	Detected				Spor	res/m3		
		<100		1K		10K	>1	100K
	Alternaria							7
	Cladosporium							107
	Total Total							114

Location: ECG-07-534A-510: 21st floor room north east area cubicle #119

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Rest	ılt: 0.3077	dF: 11 Result: 0.5932 Critical value: 0.5273 Outside Similar: Yes	Score: 101 Result: Low		
Species 3	Detected	-100		Spores/m3			
		<100	1K	10K	>100K		
Smuts, P	Cladosporium ericonia, Myxomycetes Total				27		

Location: ECG-07-534A-511: 21st floor room west area cubicle #55

% of outdoor total spores/m3	square* (indoor variation)	Agro (inc	eeme door/	nt ratio** outdoor)	corr	rman rank elation*** or/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	F	Result:	0.1538	Res Critical	dF: 12 ult: 0.3287 value: 0.4965 e Similar: No	Score: 111 Result: Low
Species]					Sp	ores/m3	
		<100		1K		10K	>100K
	Chaetomium						
	Cladosporium						80
	Total						87

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-512: 21st floor room north area cubicle #109

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		nt ratio** outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result:	0.3077	dF: 11 Result: 0.5932 Critical value: 0.5273 Outside Similar: Yes	Score: 101 Result: Low
Species 1	Detected			Spores/m3	
		<100	1K	10K	>100K
	Cladosporium				27
Smuts, P	ericonia, Myxomycetes				7
	Total				1

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

^{****} MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Client: Entek Consulting Group C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: ECG-07-534A-514: Outside ambient air, north side outside of cafeteria

Species detected		Outdoor s	ample s	pores/m3		Typical	outdo	or ranges	Freq.
	<100	1K	10K	>100K		(Nor	th An	ierica)	%
Alternaria					40	7 -	27	- 370	55
Ascospores	202 202				53	13 -	160	- 4,300	76
Basidiospores					267	13 -	320	- 14,000	93
Chaetomium					7	7 -	13	- 120	13
Cladosporium				4	1,270	53 -	530	- 8,300	95
Curvularia	0.000				20	7 -	22	- 670	16
Epicoccum	TOWNS OF THE PARTY				13	7 -	13	- 280	23
Nigrospora	1000				93	7 -	13	- 230	13
Oidium					7	7 -	13	- 210	16
Other brown					7	7 -	13	- 93	36
Penicillium/Aspergillus types					ND	27 -	210	- 2,600	87
Rusts					13	7 -	13	- 280	23
Smuts, Periconia, Myxomycetes	20				40	7 -	40	- 760	70
Ulocladium	001-4 001-3	-			20	7 -	13	- 110	6
Total				4	,850				

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: ECG-07-534A-500: 3rd floor room 322 cubicle #56

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ement ratio** oor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Re	sult: 0.3529	dF: 14 Result: 0.3901 Critical value: 0.4593 Outside Similar: No	Score: 104 Result: Low
Species	Detected			Spores/m3	
		<100	1K	10K	>100K
	Cladosporium				53
	Nigrospora				7
Penicillium/Aspergillus types		500 (5) Colored 100 (4) 100 (4)			27
Smuts, F	Smuts, Periconia, Myxomycetes				7
	Total	100			94

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-501: 3rd floor room 317 cubicle #35 at Isle Way

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rand correlation*** (indoor/outdoo	(indoor/outdoor)
Result: 2%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.5263		dF: 14 Result: 0.2879 Critical value: 0.45 Outside Similar: N	Score: 107 Result: Low
Species I	Detected			Spores/m3	-
		<100	1K	10K	>100K
	Ascospores				
	Cladosporium				27
	Curvularia				
Penici	llium/Aspergillus types				- + + + + + + + - /
Smuts, Periconia, Myxomycetes			 		27
,	Ulocladium				7
Total		dim sea	 	- 	

Location: ECG-07-534A-502: 3rd floor room 311 staging area #129

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	I	Result: 0.2667	dF: 13 Result: 0.4602 Critical value: 0.4780 Outside Similar: No	Score: 100 Result: Low	
Species 3	Detected			Spores/m3		
		<100	1K	10K	>100K	
	Cladosporium				27	
	Rusts					
	Total				34	

3800 Watt Ave., Suite 145, Sacramento, CA 95821

(650) 829-5800 Fax (650) 829-5852 www.emlab.com

Client: Entek Consulting Group C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-503: 3rd floor room 327 cubicle #94

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3750		dF: 13 Result: 0.5838 Critical value: 0.4780 Outside Similar: Yes	Score: 104 Result: Low
Species	Detected			Spores/m3	
		<100	1K	10K	>100K
	Alternaria				7
	Cladosporium				27
Smuts, Periconia, Myxomycetes					7
	Total				41

Location: ECG-07-534A-504: 11th floor room 1104 cubicle #56

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1429	dF: 13 Result: 0.6168 Critical value: 0.4780 Outside Similar: Yes	Score: 100 Result: Low
Species	Detected		Spores/m3	
		<100 IK	10K	>100K
:	Cladosporium			27
	Total			27

Location: ECG-07-534A-505: 11th floor room 1104 cubicle #32

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.2667	dF: 13 Result: 0.5920 Critical value: 0.4780 Outside Similar: Yes	Score: 103 Result: Low
Species	Detected		Spores/m3	
		<100 1K	10K	>100K
	Alternaria			
	Cladosporium			53
	Total			60

Client: Entek Consulting Group

C/O: Mr. Rick Beall Re: 07-534A; State Compensation Insurance Fund Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-506: 11th floor room 1104 cubicle #28

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1429	dF: 13 Result: 0.6168 Critical value: 0.4780 Outside Similar: Yes	Score: 100 Result: Low
Species	Detected		Spores/m3	
		<100 1K	10K	>100K
	Cladosporium	2001 100		27
	Total	V		27

Location: ECG-07-534A-507: 11th floor room 1104 cubicle #4.0

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ent ratio** /outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Resul	t: 0.1429	dF: 13 Result: 0.6168 Critical value: 0.4780 Outside Similar: Yes	Score: 100 Result: Low
Species	Detected			Spores/m3	
		<100	1K	10K	>100K
	Cladosporium				27
	Total				27

Location: ECG-07-534A-508: 21st floor room south east area cubicle #3

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)			Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 2%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.2500			dF: 14 Result: 0.3934 Critical value: 0.4593 Outside Similar: No	Score: 104 Result: Low	
Species	Detected				Spores/m3		
		<100		1K	10K	>100K	
	Ascospores					27	
Cladosporium						53	
Penicillium/Aspergillus types						27	
	Total					107	

Client: Entek Consulting Group C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-509: 21st floor room south west area cubicle #30

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 2%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.2667		dF: 13 Result: 0.5920 Critical value: 0.4780 Outside Similar: Yes	Score: 102 Result: Low	
Species 2	Detected			Spores/m3		
		<100 1K		10K	>100K	
	Alternaria				7	
	Cladosporium				107	
	Total				114	

Location: ECG-07-534A-510: 21st floor room north east area cubicle #119

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.2667		dF: 13 Result: 0.5920 Critical value: 0.4780 Outside Similar: Yes	Score: 101 Result: Low	
Species	Detected			Spores/m3		
		<100	1K	10K	>100K	
	Cladosporium				27	
Smuts, Periconia, Myxomycetes					7	
	Total				34	

Location: ECG-07-534A-511: 21st floor room west area cubicle #55

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.2667	dF: 13 Result: 0.3777 Critical value: 0.4780 Outside Similar: No	Score: 111 Result: Low	
Species	Detected		Spores/m3		
		<100 1K	10K	>100K	
	Chaetomium			7	
	Cladosporium			80	
	Total			87	

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-512: 21st floor room north area cubicle #109

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.2667		dF: 13 Result: 0.5920 Critical value: 0.4780 Outside Similar: Yes		Score: 101 Result: Low	
Species	Detected			Sp	ores/m3		
1		<100	1K		10K	>100K	
	Cladosporium					27	
Smuts, Periconia, Myxomycetes					<u> </u>	1 1 1 1 1 1 1 7	
	Total	(2003)62				34	

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

Client: Entek Consulting Group
C/O: Mr. Rick Beall
Re: 07-534A; State Compensation Insurance Fund
Date of Sampling: 09-29-2007
Date of Receipt: 10-01-2007
Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: ECG-07-534A-515: Outside ambient air, north side outside of cafeteria

o action to a minimum jum o o t		111 515. Guiside unioient un, norm bide Guiside of Guicienta								
Species detected		Outdoor	· sample s	pores/m3	3	Typical	outdo	or ranges	Freq.	
	<100	1K	10K	>100K		(Nor	th An	ierica)	%	
Alternaria	10 mg				47	7 -	27	- 370	55	
Ascospores					80	13 -	160	- 4,300	76	
Basidiospores					240	13 -	320	- 14,000	93	
Cladosporium					4,640	53 -	530	- 8,300	95	
Epicoccum					7	7 -	13	- 280	23	
Nigrospora	100				47	7 -	13	- 230	13	
Oidium					13	7 -	13	- 210	16	
Penicillium/Aspergillus types					1,430	27 -	210	- 2,600	87	
Rusts					27	7 -	13	- 280	23	
Smuts, Periconia, Myxomycetes					20	7 -	40	- 760	70	
Ulocladium					13	7 -	13	- 110	6	
Total					6,564					

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: ECG-07-534A-500: 3rd floor room 322 cubicle #56

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		nent ratio** or/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.5333		dF: 11 Result: 0.6091 Critical value: 0.5273 Outside Similar: Yes	Score: 104 Result: Low	
Species	Species Detected			Spores/m3		
		<100	1K	10K	>100K	
	Cladosporium				53	
	Nigrospora				7	
Penic	Penicillium/Aspergillus types				27	
Smuts, Periconia, Myxomycetes					7	
	Total				94	

© 2007 EMLab P&K EMLab ID: 342607, Page 1 of 6

Client: Entek Consulting Group

C/O: Mr. Rick Beall Re: 07-534A; State Compensation Insurance Fund Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-501: 3rd floor room 317 cubicle #35 at Isle Way

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Resu	lt: 0.5882	dF: 12 Result: 0.4196 Critical value: 0.4965 Outside Similar: No	Score: 107 Result: Low	
Species Detected				Spores/m3		
	-		1K	10K	>100K	
	Ascospores				27	
	Cladosporium	20.00 20.00 20.00			27	
	Curvularia				7	
Penic	illium/Aspergillus types				27	
	Smuts, Periconia, Myxomycetes				7	
Ulocladium						
	Total				102	

Location: ECG-07-534A-502: 3rd floor room 311 staging area #129

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3077		dF: 11 Result: 0.5045 Critical value: 0.5273 Outside Similar: No	Score: 100 Result: Low	
Species	Detected			Spores/m3		
		<100	1K	10K	>100K	
	Cladosporium				27	
	Rusts				7	
	Total				34	

3800 Watt Ave., Suite 145, Sacramento, CA 95821

(650) 829-5800 Fax (650) 829-5852 www.emlab.com

Client: Entek Consulting Group C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-503: 3rd floor room 327 cubicle #94

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.4286	dF: 11 Result: 0.4250 Critical value: 0.5273 Outside Similar: No	Score: 104 Result: Low	
Species 1	Detected		Spores/m3		
		<100 1K	10K	>100K	
	Alternaria				
Cladosporium				27	
Smuts, Periconia, Myxomycetes		17.00 17.00 17.00		7	
	Total			41	

Location: ECG-07-534A-504: 11th floor room 1104 cubicle #56

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1667		dF: 11 Result: 0.6295 Critical value: 0.5273 Outside Similar: Yes	Score: 101 Result: Low	
Species :	Detected			Spores/m3		
			1K	10K	>100K	
	Cladosporium				27	
	Total				27	

Location: ECG-07-534A-505: 11th floor room 1104 cubicle #32

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		cor	arman rank relation*** oor/outdoor)	MoldSCORE**** (indoor/outdoor)			
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3077		dF: 11 Result: 0.5727 Critical value: 0.5273 Outside Similar: Yes		Score: 103 Result: Low			
Species 1	Detected				S	pores/m3			
		<100		1K		10K		>100K	
	Alternaria								7
	Cladosporium								53
	Total								60

Client: Entek Consulting Group C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-506: 11th floor room 1104 cubicle #28

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1667	dF: 11 Result: 0.6295 Critical value: 0.5273 Outside Similar: Yes	Score: 101 Result: Low
Species Detected			Spores/m3	
		<100 1K	10K	>100K
Cladosporium		建		27
	Total			27

Location: ECG-07-534A-507: 11th floor room 1104 cubicle #4.0

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1667	dF: 11 Result: 0.6295 Critical value: 0.5273 Outside Similar: Yes	Score: 101 Result: Low
Species Detected			Spores/m3	
		<100 1K	10K	>100K
Cladosporium				27
	Total			27

Location: ECG-07-534A-508: 21st floor room south east area cubicle #3

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Re	esult: 0.4286	dF: 11 Result: 0.7659 Critical value: 0.5273 Outside Similar: Yes	Score: 101 Result: Low
Species Detected				Spores/m3	
		<100	1K	10K	>100K
Ascospores					27
Cladosporium					53
Penicillium/Aspergillus types					27
Total					107

Client: Entek Consulting Group C/O: Mr. Rick Beall

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

C/O: Mr. Rick Beall Date o
Re: 07-534A; State Compensation Insurance Fund Date o

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-509: 21st floor room south west area cubicle #30

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3077	dF: 11 Result: 0.5727 Critical value: 0.5273 Outside Similar: Yes	Score: 102 Result: Low
Species	Species Detected		Spores/m3	
		<100 1K	10K	>100K
	Alternaria			7
	Cladosporium			107
	Total	建		114

Location: ECG-07-534A-510: 21st floor room north east area cubicle #119

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)		
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3077				dF: 11 Result: 0.4591 Critical value: 0.52' Outside Similar: N		Score: 101 Result: Low
Species 1	Species Detected					Spores/m3		
		<100			1K	10K		>100K
Cladosporium								27
Smuts, Periconia, Myxomycetes								7
Total								34

Location: ECG-07-534A-511: 21st floor room west area cubicle #55

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.1538		dF: 12 Result: 0.3304 Critical value: 0.4965 Outside Similar: No	Score: 111 Result: Low
Species 1	Detected			Spores/m3	
		<100	1K	10K	>100K
	Chaetomium				
	Cladosporium				80
	<u>Total</u>				87

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: ECG-07-534A-512: 21st floor room north area cubicle #109

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 12 Result: 7.6813 Critical value: 21.0261 Inside Similar: Yes	Result: 0.3077	dF: 11 Result: 0.4591 Critical value: 0.5273 Outside Similar: No	Score: 101 Result: Low
Species	Detected		Spores/m3	
		<100 1K	10K	>100K
	Cladosporium			27
Smuts, F	Periconia, Myxomycetes			7
	Total	EXCERT TO THE PARTY OF THE PART		34

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.



BIOLOGICAL PARTICULATE

AIR SAMPLING RESULTS

Client: Entek Consulting Group
C/O: Mr. Rick Beall
Re: 07-534A; State Compensation Insurance Fund
Date of Sampling: 09-29-2007
Date of Receipt: 10-01-2007
Date of Report: 10-25-2007

OTHER BIOLOGICAL PARTICLES REPORT: NON-VIABLE METHODOLOGY

Location:	ECG-07	-534A-500: r room 322	ECG-07		ECG-07	-534A-502:	ECG-07	-534A-503
		cle #56	cubicle	#35 at Isle Way		area #129	3rd floor room 32 cubicle #94	
Comments (see below)	1	Vone	1	Vone	1	Vone	1	Vone
Lab ID-Version‡:	153	9501-1	153	9502-1	153	9503-1	1539504-1	
	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3
POLLEN								
Ash (Fraxinus)								
Birch (Betula)								
Cedar/Juniper (Cupressaceae)								
Chenopods (Chenopodiaceae)								
Elm (Ulmus)	2	13	2	13				
Eucalyptus (Eucalyptus)								
Grass (Poaceae)								
Mulberry (Morus)								
Oak (Quercus)				,				
Other					2	13		
Pine (Pinaceae)			-					
Ragweed (Ambrosieae)								-
Sycamore (Platanus)								
OTHER PLANT								
Algae								
Diatoms								
Fern, moss, etc. spores								
Other (wood, trichomes, etc.)	2	13	1	7	3	20	1	7
OTHER PARTICLES:								-
ANIMAL								
Epithelial (skin) cells	147	980	56	373	64	427	131	873
Hair								
Insect parts								
Mites								
FUNGI								
Hyphal fragments	2	13	1	7	2	13	1	7
NON-BIOLOGICAL								
Cotton fibers							2	13
Glass fiber								
Soot								
Background debris (1-4+)†	1+		1+		1+		1+	
Sample volume (liters)	150		150		150		150	

Comments:

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[†] Background debris is an indication of the amounts of non-biological particulate matter present on the slide (dust in the air) and is graded from 1+ to 4+ with 4+ indicating the largest amounts. To evaluate dust levels it is important to account for differences in sample volume.

Client: Entek Consulting Group
C/O: Mr. Rick Beall
Re: 07-534A; State Compensation Insurance Fund
Date of Sampling: 09-29-2007
Date of Receipt: 10-01-2007
Date of Report: 10-25-2007

OTHER BIOLOGICAL PARTICLES REPORT: NON-VIABLE METHODOLOGY

Location:	11th f	-534A-504: oor room ubicle #56	11th fl	7-534A-505: oor room ubicle #32	11th fl	-534A-506: oor room ubicle #28	11th f	'-534A-507: loor room ubicle #4.0
Comments (see below)	1	Vone	Ŋ	Vone	Ŋ	Vone	1	Vone
Lab ID-Version‡:	153	9505-1	153	9506-1	153	9507-1	153	9508-1
	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3
POLLEN								
Alder (Alnus)								
Ash (Fraxinus)								
Birch (Betula)								
Cedar/Juniper (Cupressaceae)								
Chenopods (Chenopodiaceae)								
Elm (Ulmus)	1	7						
Eucalyptus (Eucalyptus)								
Grass (Poaceae)								
Mulberry (Morus)					-			
Oak (Quercus)								
Other								
Pine (Pinaceae)								
Ragweed (Ambrosieae)								
Sycamore (Platanus)	1							
OTHER PLANT								
Algae			***************************************					
Diatoms				-				
Fern, moss, etc. spores								
Other (wood, trichomes, etc.)	2	13	2	13			1	7
OTHER PARTICLES:			****					
ANIMAL								
Epithelial (skin) cells	121	807	142	947	101	673	73	487
Hair					,			
Insect parts								
Mites					•			
FUNGI								
Hyphal fragments	1	7	1	7	11	7		
NON-BIOLOGICAL								
Cotton fibers	2	13			3	20	3	20
Glass fiber	1	7	11	7	1	7		
Soot								
Background debris (1-4+)†	1+		1+		1+		1+	
Sample volume (liters)	150		150		150		150	

Comments:

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[†] Background debris is an indication of the amounts of non-biological particulate matter present on the slide (dust in the air) and is graded from 1+ to 4+ with 4+ indicating the largest amounts. To evaluate dust levels it is important to account for differences in sample volume.

Client: Entek Consulting Group
C/O: Mr. Rick Beall
Re: 07-534A; State Compensation Insurance Fund
Date of Sampling: 09-29-2007
Date of Receipt: 10-01-2007
Date of Report: 10-25-2007

OTHER BIOLOGICAL PARTICLES REPORT: NON-VIABLE METHODOLOGY

Location:	21st fl south cub	-534A-508: oor room east area sicle #3	21st fl south cubi	-534A-509: oor room west area icle #30	om 21st floor room north east area		ECG-07-534A-511: 21st floor room west area cubicle # 55	
Comments (see below)	1	Vone	1	None	N	None	1	Vone
Lab ID-Version‡:	153	9509-1	1539510-1		153	9511-1	1539512-1	
	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3
POLLEN								
Ash (Fraxinus)								
Birch (Betula)								
Cedar/Juniper (Cupressaceae)								
Chenopods (Chenopodiaceae)								
Elm (Úlmus)							1	7
Eucalyptus (Eucalyptus)								
Grass (Poaceae)					1			
Mulberry (Morus)			÷					
Oak (Quercus)								
Other								
Pine (Pinaceae)								
Ragweed (Ambrosieae)								
Sycamore (Platanus)								
OTHER PLANT								
Algae			·					
Diatoms								
Fern, moss, etc. spores								
Other (wood, trichomes, etc.)	2	13			1	7	1	7
OTHER PARTICLES:								
ANIMAL								
Epithelial (skin) cells	110	733	88	587	67	447	54	360
Hair								
Insect parts								
Mites			·				:	
FUNGI								
Hyphal fragments								
NON-BIOLOGICAL								
Cotton fibers	4	27	3	20	11	7	3	20
Glass fiber			1	7				
Soot			×'					
Background debris (1-4+)†	1+		1+		1+		1+	
Sample volume (liters)	150		150		150		150	

Comments:

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[†] Background debris is an indication of the amounts of non-biological particulate matter present on the slide (dust in the air) and is graded from 1+ to 4+ with 4+ indicating the largest amounts. To evaluate dust levels it is important to account for differences in sample volume.

Client: Entek Consulting Group
C/O: Mr. Rick Beall
Re: 07-534A; State Compensation Insurance Fund
Date of Sampling: 09-29-2007
Date of Receipt: 10-01-2007
Date of Report: 10-25-2007

OTHER BIOLOGICAL PARTICLES REPORT: NON-VIABLE METHODOLOGY

Location:	21st fl north are	-534A-512: oor room ea cubicle # 109	Outside a north sid	-534A-513: ambient air, e outside of feteria	Outside a	ECG-07-534A-514: Outside ambient air, north side outside of cafeteria		ECG-07-534A-515: Outside ambient air, north side outside of cafeteria	
Comments (see below)	1	Vone	N	None	N	Vone	None		
Lab ID-Version‡:	153	9513-1	153	9514-1	153	9515-1	153	9516-1	
	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3	raw ct.	particles/m3	
POLLEN									
Birch (Betula)									
Cedar/Juniper (Cupressaceae)									
Chenopods (Chenopodiaceae)									
Elm (Úlmus)			8	53	4	27	2	13	
Eucalyptus (Eucalyptus)									
Grass (Poaceae)									
Mulberry (Morus)									
Oak (Quercus)		-							
Other			2	13	1	7			
Pine (Pinaceae)					1	7			
Ragweed (Ambrosieae)									
Sycamore (Platanus)									
OTHER PLANT								:	
Algae									
Diatoms									
Fern, moss, etc. spores									
Other (wood, trichomes, etc.)	1	7	22	147	19	127	11	73	
OTHER PARTICLES:	100								
ANIMAL									
Epithelial (skin) cells	54	360	17	113	34	227	16	107	
Hair									
Insect parts									
Mites									
FUNGI									
Hyphal fragments			11	73	8	53	2	13	
NON-BIOLOGICAL									
Cotton fibers	2	13	1	7	2	13	3	20	
Glass fiber	1_1_	7	1	7	2	13	4	27	
Soot			6	40	7	47	9	60	
Background debris (1-4+)†	1+		2+		2+		2+		
Sample volume (liters)	150		150		150		150		

Comments:

The Limit of Detection and Minimum Reporting Limit is a raw count of 1. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[†] Background debris is an indication of the amounts of non-biological particulate matter present on the slide (dust in the air) and is graded from 1+ to 4+ with 4+ indicating the largest amounts. To evaluate dust levels it is important to account for differences in sample volume.



BULK SAMPLING RESULTS CARPET DUST CULTURE FOR MOLD SPORES



BULK MATERIAL Analysis Report Form for ENTEK CONSULTING GROUP, INC.

4200 Rocklin Road, Suite 7 Rocklin, CA 95677 (916) 632-6800 (916) 632-6812 Fax

Date of Sampling: 09/29/07

Lab:

EML - San Bruno

Job Number: 07-534A

Analysis Requested:

Biological Evaluation -

Culture

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization

Turnaround Time:

Standard

450 N Street

Sacramento, CA

SAMPLE#	RESULTS	MATERIAL DESCRIPTION/LOCATION
ECG-07-534A-600	SEE ATTACHED LABORATORY REPORT	Carpet Dust Sample From Room 2217 Law Library; 22 nd Floor; Approximately 4 Square Feet
ECG-07-534A-601	SEE ATTACHED LABORATORY REPORT	Carpet Dust Sample From Open Office Area Near Room 2206 and Cubicle #095

Samples Collected on to 0.45µ MCE Filters in 25mm Cassettes at a Flow of Approximately 12-15 LPM.

EMLab ID: 342865, Page 1 of 1

1150 Bayhill Drive, Suite 100, San Bruno, CA 94066 (650) 829-5800 Fax (650) 829-5852 www.emlab.com

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-02-2007 Date of Report: 10-11-2007

FUNGAL CULTURE REPORT

Location:	Carpet dust sar library, 22nd	ECG-07-534A-600: ECG-07-534A Carpet dust sample from room 2217 law library, 22nd floor, approximately 4 square feet ECG-07-534A			-601: n open office l cubicle #095		
Comments (see below)		None		None			
Sample type]	Dust samp	le		Dust sample	9	
Media used	Cellu	lose/DG18	/MEA	Cellu	lose/DG18/		
Lab ID-Version‡:		1500991-	l		1500992-1		
	sample ct.†	%	cfu*/unit	sample ct.†	%	cfu*/unit	
Acremonium							
Alternaria	200	3	8,000	20	6	800	
Aspergillus flavus						000	
Aspergillus fumigatus							
Aspergillus nidulans							
Aspergillus niger	100	2	4,000				
Aspergillus ochraceus							
Aspergillus versicolor	400	7	16,000			****	
Aureobasidium		-					
Bipolaris/Drechslera group							
Botrytis							
Cladosporium	2,800	45	110,000	160	50	6,400	
Curvularia	100	2	4,000	100		0,400	
Epicoccum	200	3	8,000	30	9	1,200	
Fusarium	100	2	4,000			1,200	
Non-sporulating fungi				20	6	800	
Paecilomyces						000	
Penicillium	1,200	20	48,000	80	25	3,200	
Rhizopus				10	3	400	
Stachybotrys chartarum				10	-	700	
Ulocladium							
Yeasts	1,100	18	44,000				
Dilutions††	1:10, 1:100, 1:1,000	& 1:10,000	,,,,,,,	1:10, 1:100, 1:1,000	& 1:10,000		
Sample size	0.025			0.025			
Unit	1 gram			1 gram			
TOTAL CFU*/unit * cfu = colony forming units			246,000	1 514111		12,800	

* cfu = colony forming units

Caution should be used when interpreting percentages. Totals may not equal 100 due to rounding.

Comments:

†Sample count is the calculated number of colonies that would have grown if the entire selected sample size analyzed were plated out. ††Results represent a compiled result from multiple media and multiple dilutions. Sensitivity of the results depends largely upon the dilutions used and the size of the sample. For example, a dilution of 1:100 means that 1 colony on a plate represents a sample count of 100. For a sample of 0.025 grams, this would represent 4,000 cfu/gram. For a sample of 0.002 grams, this would represent 50,000 cfu/gram. When detected, the minimum detection and reporting limit is a colony count of 1 at the lowest dilution plated. Interpretation is left to the company and/or persons who conducted the field work. ‡ A "Version" greater than 1 indicates amended data.



BULK SAMPLING RESULTS FOR MOLD BY

DIRECT MOUNT MICROSCOPIC EVALUATION



BULK MATERIAL Analysis Report Form

ENTEK CONSULTING GROUP, INC. 4200 Rocklin Road, Suite 7 Rocklin, CA 95677 (916) 632-6800 (916) 632-6812 FAX

Date of Sampling: 09/29/07

Lab: EML - Sacramento

Job Number: 07-534A

Analysis Requested: Mold by Direct Mount

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization

Turnaround Time:

Standard

450 N Street

Sacramento, CA

SAMPLE#	RESULTS	MATERIAL DESCRIPTION/LOCATION
ECG-07-534A-100	SEE ATTACHED LABORATORY REPORT	South Wall Base Wall Near South East Corner - 22 nd Floor Across From Cubicle 095 - Suspect Black Mold
ECG-07-534A-101	SEE ATTACHED LABORATORY REPORT	Base of West Wall Near Door of Room 2238 on Floor 22; Suspect Brown Behind Base Cove
ECG-07-534A-102	SEE ATTACHED LABORATORY REPORT	South Wall of Cubicle East of #33 at Opening Room 2305; 23 rd Floor; Suspect Brown Mold

Z:\Clients\State Comp Ins Fund\07-534A Board of Equal 450 N Street\Bikother.Rpt100.wpd

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

DIRECT MICROSCOPIC EXAMINATION REPORT

(Wet Mount)

Debris and/or S Description	Miscellaneous Spores Present*	MOLD GROWTH: Molds seen with underlying mycelial and/or sporulating structures†	Other Comments††	General Impression						
1 ,	ipe sample ECG-07-534A-10									
1 ,		Lab ID-Version‡: 1499761-1: Tape sample ECG-07-534A-100: South wall base wall near south east corner, 22nd floor across from cubicle 095, suspect black mold								
	Very few	4+ Stachybotrys species 2+ Acremonium species	None	Mold growth						
Lab ID-Version: 1499762-1: T	Tane sample ECG 07 52	4A 101: Boss of want will be a consideration								
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	rape sample ECO-07-53	4A-101: Base of west wall near door of room 2238 or	ı floor 22, suspect brown be	hind base cove						
Heavy	Very few	1+ colorless spores typical of Penicillium / Aspergillus	None	Mold growth						
I ah ID V 1400762 1 7										
Lau 1D-version: 1499/63-1: T	Tape sample ECG-07-53	4A-102: South wall of cubicle east of #33 at opening	room 2305, 23rd floor, susp	ect brown mold						
Light	1	4+ Aspergillus species	None	Mold growth						
* Indicative of normal condi										

^{*} Indicative of normal conditions, i.e. seen on surfaces everywhere. Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating. Distribution of spore types seen mirrors that usually seen outdoors.

[†] Quantities of molds seen growing are listed in the MOLD GROWTH column and are graded 1+ to 4+, with 4+ denoting the highest numbers.

^{††} Some comments may refer to the following: Most surfaces collect a mix of spores which are normally present in the outdoor environment. At growth. Marker genera are those spore types which are present normally in very small numbers, but which multiply indoors when conditions are favorable for growth.

[‡] A "Version" greater than 1 indicates amended data.

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

DIRECT MICROSCOPIC EXAMINATION REPORT

DIRECT MICROSCOTIC EX	AMINATION REPORT		
Location:	ECG-07-534A-100:	ECG-07-534A-101:	ECG-07-534A-102:
	South wall base wall near	Base of west wall near	South wall of cubicle east
	south east corner, 22nd	door of room 2238 on	of #33 at opening room
	floor across from cubicle	floor 22, suspect brown	2305, 23rd floor, suspect
Carrata	095, suspect black mold	behind base cove	brown mold
Sample type:	Tape sample	Tape sample	Tape sample
Lab ID-Version‡:	1499761-1	1499762-1	1499763-1
MOLD/FUNGAL GROWT	H*: Molds seen growing with	underlying mycelial and/or spe	orulating structures
Acremonium	2+		
Alternaria			
Aspergillus			4+
Aureobasidium			
Basidiospores			
Chaetomium			
Cladosporium			
Colorless spores typical of Penicillium / Aspergillus		1+	
Fusarium			
Other colorless, ID unknown			
Stachybotrys	4+		
Torula			
Ulocladium			
Miscellaneous spores**	Very few	Very few	Very few
Other comments†	None	None	None
Background debris or Description††	Light	Heavy	Light
General impression	Mold growth	Mold growth	Mold growth

Interpretation is left to the company and/or persons who conducted the field work.

‡ A "Version" greater than 1 indicates amended data.

^{*} See Mold/Fungal Growth Details table on the last page.

^{**} See Miscellaneous Spores table on the last page.

[†] Some comments may refer to the following: Most surfaces collect a mix of spores which are normally present in the outdoor environment. At times it is possible to note a skewing of the distribution of spore types, and also to note "marker" genera which may indicate indoor mold growth. Marker genera are those spore types which are present normally in very small numbers, but which multiply indoors when conditions are favorable for growth.

^{††} Background debris is an indication of the amounts of non biological particulate matter present. This background amorphous material is graded and described as scant, light, moderate, heavy, or very heavy. (Very heavy background debris may obscure visibility.)

Client: Entek Consulting Group

C/O: Mr. Rick Beall

Re: 07-534A; State Compensation Insurance Fund

Date of Sampling: 09-29-2007 Date of Receipt: 10-01-2007 Date of Report: 10-02-2007

Mold/Fungal Growth Rating Details

gar Growth Rading Details						
Quantities of molds indicating growth are listed in the MOLD/FUNGAL GROWTH section. Judgement is used in determining the amount of growth present in the sample. For example, if only one portion of the sample has evidence of heavy growth, then it will receive a rating of heavy growth even though, strictly speaking, on a percentage basis of the entire sample, the amount of growth is low.						
Swab/Tape/Dust/Wipe sample	Bulk Sample					
Evidence of very light growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in less than 10% of the microscopic fields examined.	Areas of very light growth detected by the presence of spores of one type seen with underlying mycelial and/ or with their sporulating structures in the bulk sample.					
Evidence of light growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 10 to 25% of the microscopic fields examined.	Areas of light growth detected by the presence of spores of one type seen with underlying mycelial and/or with their sporulating structures in the bulk sample.					
Evidence of moderate growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 26 to 50% of the microscopic fields examined.	Areas of moderate growth detected by the presence of spores of one type seen with underlying mycelial and/ or with their sporulating structures in the bulk sample.					
Evidence of heavy growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 51 to 75% of the microscopic fields examined.	Areas of heavy growth detected by the presence of spores of one type seen with underlying mycelial and/ or with their sporulating structures in the bulk sample.					
Evidence of very heavy growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found to be nearly confluent in the majority of the microscopic fields examined.	Areas of very heavy growth detected by the presence of spores of one type seen with underlying mycelial and/ or with their sporulating structures in the bulk sample.					
	Quantities of molds indicating growth are listed in the N in determining the amount of growth present in the san evidence of heavy growth, then it will receive a rating percentage basis of the entire san Swab/Tape/Dust/Wipe sample Evidence of very light growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in less than 10% of the microscopic fields examined. Evidence of light growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 10 to 25% of the microscopic fields examined. Evidence of moderate growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 26 to 50% of the microscopic fields examined. Evidence of heavy growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 51 to 75% of the microscopic fields examined. Evidence of very heavy growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 51 to 75% of the microscopic fields examined.					

Miscellaneous Spores

Slides/specimens are examined for the presence of mold spores and pollen, noting the quantities and distribution of spore types found. A designation of 'normal trapping' is made when a mix of spore types is present with the same general distribution as is usually found outdoors. In other words, the biological component of the sample surface is like that found everywhere. Types of spores present would include basidiospores (mushroom spores), myxomycetes (slime molds), plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating. Many of these spore types would not be found growing indoors on building materials since many plant pathogens require living plants for growth, and mushrooms require compost, leaf duff of various types, or associations with roots of certain trees, etc. Due to these factors, when a mix of spores seen include these types as well as pollen, the rational source is the outside air, rather than indoor mold growth. The numbers of miscellaneous spores seen are graded and described as shown below as none, very few, few, variety, and wide variety.

None	Very Few	Few	Variety	Wide Variety
No spores detected	Very few spores detected	A few spores detected	Many spores containing a variety of different genera detected	Many spores containing a wide variety of different genera detected



BIOLOGICAL PARTICULATE

SETTLED DUST SAMPLING RESULTS

BY

POLARIZED LIGHT MICROSCOPY



BULK MATERIAL Analysis Report Form for ENTEK CONSULTING GROUP, INC.

4200 Rocklin Road, Suite 7 Rocklin, CA 95677 (916) 632-6800 (916) 632-6812 Fax

Date of Sampling: 09/29/07

Lab: Forensic Analytical Specialties, Inc.

Job Number: 07-534A

Analysis Requested: Particle ID Analysis

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization

Turnaround Time:

Standard

450 N Street Sacramento, CA

SAMPLE#	RESULTS	MATERIAL DESCRIPTION/LOCATION
ECG-07-534A-700	SEE ATTACHED LABORATORY REPORT	Settled Dust / Particulate on Honeywell HEPA Filter Unit at Judy Knights Office in Room 317 at Cubicle #44

BULK MATERIAL Analysis Request Form for ENTEK CONSULTING GROUP, INC.

4200 Rocklin Road, Suite 7 Rocklin, CA 95677 (916) 632-6800 (916) 632-6812 Fax

Date of Sampling: 09/29/07

Lab:

Forensic Analytical Specialties, Inc.

Job Number: 07-534A

Analysis Requested: Particle ID Analysis

Client Name: State Compensation Insurance Fund

Collected by:

Rick Beall

Site Address: Board of Equalization

Turnaround Time:

Standard

450 N Street

Sacramento, CA

Special Instructions: Attn: Mark Floyd

SAMPLE#	MATERIAL DESCRIPTION/LOCATION
ECG-07-534A-700	Settled Dust / Particulate on Honeywell HEPA Filter Unit at Judy Knights Office in Room 317 at Cubicle #44

Delivered by:	MUK	VIA FEDER	Date:_	10-1-07	Time: /:/5PM
Received by:	AJG		Date:	(O) RE	CEI VIM eOCT 0 2 2007



3777 Depot Road, Suite 409, Hayward, California 94545 Phone: 510-887-8828, Fax: 510-887-4218

PARTICLE IDENTIFICATION ANALYSIS by Polarized Light Microscopy (PLM)

Entek Consulting Group Rick Beall

4200 Rocklin Rd, Suite 7 Rocklin CA 95677

Page: Client Number: 1/1 A31353

Report Number: SP Number:

T013590 07109

Date Received:

10/2/07

Site:

Board of Equalization, Sacramento

Date Reported:

6/14/07

Job#	b# 07-534A, State Fund Compensation Insurance		Analyst:	LW	
		PARTICLE IDENTIFICATION ANALYSIS RES	BULTS		
Client Sample No.		ECG-07-534A-700	MT.		
Lab Sam	ple No.	20052294			
Description:		Settled dust/Particulate on Honeywell HEPA Filter Unit			
		at Judy Knight's office, Room 317, Cubicle #44			
Fibrous	Major	Cotton			
		Cellulose			
	Minor	Paper			
	Trace	Synthetics			
		Wool			
		Mineral wool			
		Trichomes			
		Hair, feline			
Non-	Major	Epithelial cells			
Fibrous	Minor	Clear isotropics*			

i ibious	IVIIIIOI	Clear isotropics
	Trace	Pollen: pinaceae, poaceae
		Iron oxide
		Limestone
		Feldspars
		Opaques
		Organic debris
		Paint chips
		Quartz
		Spray paint
		Starch, agglomerated
		Salt (NaCl)

Quantitation: Major: >10%, minor: 1-10%, trace: <1%.

Phenolic foam

^{*} Clear isotropics may be glass chips or aluminum corrosion products

